

# Stock Quality Assessment: Forecasting Survival or Performance on a Reforestation Site

Steven C. Grossnickle and Raymond S. Folk

*Research scientists, Forest Biotechnology Centre  
B.C. Research Inc., Vancouver, British Columbia*

*Seedlings are exposed to a wide range of environmental conditions on reforestation sites. These conditions could result in stress that reduces survival and/or growth of newly planted seedlings. Field survival and field performance potential need to be distinct areas of evaluation when selecting and interpreting stock quality tests. Tests that measure the functional integrity of seedlings help forecast their survival capability. Tests that simulate anticipated field environmental conditions help forecast a seedling's physiological performance and potential for growth on a reforestation site. Tree Planters' Notes 44(3): 113-121; 1993.*

Stock quality assessment has evolved to include both morphological and physiological tests (see reviews by Sutton 1979, Chavasse 1980, Jaramillo 1980, Schmidt-Vogt 1981, Ritchie 1984, Duryea 1985a, Glerum 1988, Lavender 1988, Puttonen 1989, Hawkins and Binder 1990, Johnson and Cline 1991, Omi 1991). The wide array of testing procedures has sometimes led to confusion in selection of tests for specific purposes. Part of this confusion stems from the fact that stock quality tests can have one of two different purposes: evaluating nursery development (for example, determining nursery growth phase or evaluating readiness for lifting and storage) or forecasting field survival and/or growth (Duryea 1985b). A clearer understanding of the nature and purpose of specific testing techniques will help nursery personnel and regeneration silviculturists choose appropriate tests and make more effective decisions.

With any type of stock quality assessment procedure, differences in test results could be due to species, genetic variability of seedlots, variations in nursery culture, cold or frozen storage regimes, and variations in testing conditions. Separate testing standards need to be developed for seedlings produced from various combinations of the above nursery decisions. Seedling users also need to be aware that the mishandling of stock during transport to planting sites, improper planting procedures, and unpredictability of field site environmental conditions will influence how test results agree with initial seedling survival and/or growth on a reforestation site.

Stock quality assessment as it relates to forecasting either initial field survival or field performance potential (i.e., potential for initial growth on a reforestation site) is the focus of this paper. Testing procedures are discussed and evaluated for their suitability to provide information on these aspects of stock quality assessment. Understanding the benefits and limitations of these testing approaches will provide nursery personnel and regeneration silviculturists with a better appreciation of their potential utility within an operational forest regeneration program.

## **Planting Stress and Stock Quality Assessment**

Seedlings can be exposed to stress just after they are planted on a reforestation site. This is usually attributable to water stress because root confinement, poor contact of roots with soil, and low root system permeability can limit water uptake from the soil needed to meet transpirational demands placed upon seedling shoot systems by atmospheric conditions (Kozlowski and Davies 1975, Burdett 1990). Planting stress will be overcome only if seedlings have functional physiological processes required for morphological development, primarily root growth, to occur. When root growth occurs in newly planted seedlings, water stress is reduced and a seedling's physiological processes then have the capability to respond in a normal manner (Sands 1984, Grossnickle 1988, Carlson and Miller 1990, Brissette and Chambers 1992).

Further limitations on the physiological processes of newly planted seedlings can occur from exposure to environmental extremes on a reforestation site. The most dramatic of these are alterations in heat exchange processes and site-water relations (Miller 1983). Low temperature and drought conditions are two predominant types of environmental stress occurring on reforestation sites.

First, freezing events can cause frost damage (Nilsson and Eriksson 1986, Grossnickle et al. 1991b) and/or reduced gas exchange capability (Neilson and Jarvis 1976, DeLucia 1987, Grossnickle and Arnott 1992) in newly planted seedlings. Low soil temperature condi-

tions in early spring can cause reduced root growth (Nambiar et al. 1979, Lopushinsky and Kaufmann 1984, Grossnickle et al. 1991b), and/or restrict water uptake, resulting in water stress (Kaufmann 1977, Nambiar et al. 1979, Lopushinsky and Kaufmann 1984, Grossnickle 1988).

Second, newly planted seedlings can be exposed to drought through limited soil moisture and/or high evaporative demand conditions of the atmosphere. Drought conditions cause seedling water stress by restricting water uptake from the soil (Kaufmann 1979, Dixon et al. 1983, Grossnickle and Reid 1984, Sands 1984, Livingston and Black 1987a, Brissette and Chambers 1992) and by inadequate stomatal control as evaporative demand increases (Grossnickle and Blake 1987, Livingston and Black 1987b, Grossnickle and Arnott 1992). The result of increased water stress in newly planted seedlings is a reduction in growth (Nambiar and Zed 1980, Margolis and Waring 1986, Livingston and Black 1988, Grossnickle and Heikuri-nen 1989). As a result, planting stress can be exacerbated by field site environmental conditions that reduce growth and delay a seedling's capability to occupy the site.

No stock quality assessment program can alleviate the stress seedlings are exposed to on reforestation sites. However, a program that defines a seedling's functional integrity could determine whether it has the capability to survive potentially stressful environmental conditions, because initial field survival is dependent on whether a seedling has the physiological capability to function normally at time of planting. On the other hand, a program that defines field performance potential by measuring a seedling's physiological responses and morphological development under simulated environmental conditions of the planting site would provide information on field growth potential. Though testing for field performance potential would provide information on survival capability, there is no guarantee that testing for survival would provide sufficient information on field performance potential. Thus, stock quality assessment as it relates to a seedling's initial field survival or field performance potential are considered distinct areas of evaluation and are examined as separate topics.

### Field Survival Capability

Currently, there are a number of testing procedures that provide information on the initial survival potential of operationally produced stock. These tests measure a seedling's vitality under a specific set of conditions that defines a certain level of quality when

tested (Ritchie and Tanaka 1990, Langerud 1991). These kinds of tests measure the functional integrity of seedlings, which helps determine their initial survival capability. Functional integrity indicates whether a seedling is, or is not, damaged to the point of limiting primary physiological processes. The intent of these testing approaches is to remove seedlings that do not meet certain minimum physiological performance standards (i.e., the "bad apple concept"). Seedlings that meet minimum standards probably have a greater capability to survive in all but the most severe of field site environmental conditions (Sutton 1988).

The following are examples of testing procedures that provide information on the functional integrity of tested seedlings. These tests have been developed for the purpose of batch-culling poorly grown and handled seedlings. They are used to categorize large groups of seedlings, all having a similar nursery culture regime or from a similar seed source, by measuring a subsample from the entire population. A brief description of each test is given below. Further specific information on each testing procedure can be found in the cited articles.

1. **Root growth capacity** is a measure of a seedling's ability to regenerate new roots and an indirect measure of a seedling's overall physiological condition (Stone 1955, Ritchie and Dunlap 1980, Ritchie 1985, Burdett 1987, Ritchie and Tanaka 1990, Sutton 1990).
2. **Oregon State University vigor test** is a measure of a seedling's subsequent survival after exposure to a single controlled stress event (15 minutes at 30 °C and 30% relative humidity) (McCreary and Duryea 1985, 1987; Lavender 1988).
3. **Shoot water potential** of potted seedlings after a set time period is an indirect measure of a root system's capability to absorb water and thus maintain a proper seedling water balance (McCreary and Duryea 1987).
4. **Needle conductance** (Orlander and Rosvall-Ahnebrink 1987) and **transpiration** (Langerud et al. 1991) are measures of the water movement capability of needles and an indirect measure of a root system's capability to absorb water and the xylem's capacity to transport water to the needles.
5. **Infrared thermography** is a measure of foliage heat exchange (i.e., temperature) resulting from transpiration and an indirect measure of a root system's capability to absorb water and the xylem's capability to transport water to the needles (Weatherspoon and Laacke 1985, Orlander et al. 1989).

6. **Root system water loss capability** measured under positive pressure is an indirect measure of root system integrity (Ritchie 1990).
- 7 **Fine root electrolyte leakage** is an indirect measure of root system integrity (McKay and Mason 1991, McKay 1992).
8. **Variable chlorophyll fluorescence** is a measure of photosynthesis and an indirect measure of a seedling's overall physiological condition (Vidivar et al. 1989, 1991).
9. **Stress-induced volatile emissions** is a measure of cell injury due to membrane breakdown (Hawkins and DeYoe 1992).

The above tests measure different morphological or physiological parameters in relation to initial field survival of tested seedlings. Seedlings that do not meet certain minimum performance standards usually have poor field survival capability. On the other hand, seedlings that meet certain minimum performance standards have a greater capability to survive under typical reforestation site conditions.

However, no single testing procedure accurately forecasts field survival under all circumstances. For example, an extensive operational test of root growth capacity (RGC) found that RGC had a poor relationship with field survival under some circumstances (Binder et al. 1988). Seedlings with poor RGC had a higher probability of increased mortality. However, they found that even seedlings with high RGC could still have an unacceptable mortality level after field planting. This example emphasizes the limitations inherent in using a single test as an indicator of a seedling's overall quality. Seedlings have a wide array of physiological processes that continually respond to environmental conditions. Proper stock quality assessment must consider the dynamic and interdependent nature of a seedling's physiological processes.

### Field Performance Potential

A seedling's performance on a reforestation site depends on its inherent growth potential and the degree to which the environmental conditions of the field site allow this growth potential to be expressed. Thus, the degree to which a seedling can adapt to site conditions just after planting influences its initial growth on the reforestation site (Burdett 1983). To determine a seedling's field performance potential, the seedling should be assessed in relation to anticipated environmental conditions at the site (Duryea 1985b; Sutton 1982, 1988; Puttonen 1989; Grossnickle et al. 1988, 1991a; Hawkins and Binder 1990). In addition, an array of morphological and physiological tests

that examine factors important for determining a seedling's field performance potential is required because stock quality reflects the expression of a multitude of physiological and morphological attributes (Ritchie 1984). An array of tests that simulate anticipated field environmental conditions would help forecast seedlings physiological performance and potential for growth on a reforestation site.

To measure a seedling's physiological response and growth under a range of environmental conditions, tests should define performance under optimum environmental conditions, as well as define stress tolerance and avoidance parameters (Levitt 1980). This approach was first presented by Timmis (1980), who developed a series of tests to simulate essential physiological responses and growth behavior of seedlings in any environment and derived numerical values for these responses. Examples of possible material and performance attribute tests important in defining a seedling's field performance potential are shown in table 1. In tests measuring performance attributes, whole seedlings are subjected to some test condition that integrates their response over time or to a range of environmental conditions (Ritchie 1984). In tests measuring material attributes, an individual morphological or physiological parameter of the seedling is tested (Ritchie 1984).

Seedlings are normally exposed to some type of stress after planting on a reforestation site. Anticipated environmental conditions could be defined by reforestation silviculturists during on-site development of regeneration prescriptions. Test environments could then be selected that match this range and combination of anticipated environmental conditions.

Effective determination of field performance potential depends on the selection of a smaller number of morphological and physiological attributes from a master table (table 1). As described earlier, low temperature and drought are two predominant types of environmental stress that could occur on reforestation sites. Possible attributes to consider measuring on seedlings to be planted on potentially cold or droughty reforestation sites are described in figures 1 and 2, respectively. This approach to stock quality assessment is designed to allow the user to have information from a number of material and performance attribute tests that are important for their intended purpose.

Results from testing programs could be integrated to develop a means of expressing the overall physiological and morphological quality of seedlings. The performance potential index (PPI) has been developed to integrate material and performance attribute tests for a comprehensive perspective of seedling field per-

**Table 1** -Possible material (morphological and physiological) and performance attribute tests and their intended purposes for defining field performance potential

#### Morphological attribute tests

**Height:** General measure of photosynthetic capacity and transpirational area (Armson and Sadreka 1979); greater height is an advantage on sites where brush competition and animal browsing are potential problems (Cleary et al. 1978).

**Diameter:** General measure of a seedling's durability, root system size, and protection from drought and heat damage; provides support to withstand physical abuse (Cleary et al. 1978).

**Needle surface area:** Direct measure of potential photosynthetic or transpirational surface area.

**Root surface area or dry weight:** Good indicator of absorptive root surface (Thompson 1985).

**Needle primordia:** Important indicator of shoot growth potential (Colombo 1986).

**Seedling water balance ratio (needle dry weight/[stem diameter x root dry weight]):** Measure of drought avoidance potential for situations where water absorption lags behind transpiration (Grossnickle et al. 1991 a).

#### Physical attribute tests

**Osmotic potential at turgor loss point:** Quantitative measure of drought tolerance (Tyree and Jarvis 1982).

**Maximum bulk modulus of elasticity:** Quantitative measure of cells' elasticity, with greater elasticity representing greater turgor maintenance (Tyree and Jarvis 1982).

**Seedling water movement:** Measurement of water movement capability in relation to a plant's resistances along the pathway (i.e., root xylem, needle) to the atmosphere (Hinckley et al. 1978); provides measure of drought avoidance potential.

**Cuticular transpiration:** Measure of needle's capability to avoid water loss after stomata have theoretically closed (Vanhinsberg and Colombo 1990).

**Days to terminal budbreak:** Direct measure of bud dormancy status (Lavender 1991) and indirect measure of changes in drought and cold temperature tolerance (Burr 1990).

#### Performance attribute tests

**Root growth capacity:** General indicator that all systems in a seedling are functioning properly (Ritchie 1984) and measure of seedling performance potential (Burdett 1987).

**Root growth capacity at low root temperature or after exposure to drought conditions:** Measure of a seedling's performance and root growth capability under stressful soil conditions (Grossnickle et al. 1991 a).

**Frost hardiness:** Measure of a seedling's tolerance to freezing temperatures (Glerum 1985).

**Net photosynthesis 14-day integral under optimal environmental conditions:** Direct measure of a seedling's photosynthetic capability (Grossnickle et al. 1991 a).

**Net photosynthesis 14-day integral at low root temperatures:** Direct measure of seedling tolerance to low temperatures (Grossnickle et al. 1991 a).


**Net photosynthetic capability at decreasing predawn water potentials:** Direct measure of a seedling's tolerance to drought (Grossnickle et al. 1991 a).

**Gas exchange capability at various vapor pressure deficits:** Measure of stomatal conductance, transpiration, and/or net photosynthesis used to define the efficiency of a plant's CO<sub>2</sub> uptake in relation to water loss (Landsberg 1986).

formance potential (Grossnickle et al. 1991c). The PPI provides a means for collectively interpreting the results from a group of tests within a standardized, yet quantitative framework. The PPI, measured immediately before planting, has been used to clarify the relationship between nursery culture regimes (Grossnickle et al. 1991a-c) or stock types (Grossnickle and Major 1993a,b) with field performance. Another approach to integrating test results has been proposed by D'Aoust et al. (1991). Their approach characterizes seedling performance potential with ten morphological and physiological parameters. Principal component analysis was used to identify a smaller set of parameters that adequately represent information contained in the whole set. Measurement of four variables (i.e., diameter, stem height, shoot water potential at planting, and root growth capacity) before field planting were sufficient to characterize the morphology and physiology of the seedlings produced.

However, limitations are inherent in stock quality assessment depending on when the test is used and what morphological and physiological attributes of the seedlings are measured (Puttonen 1989). These limitations influence the usage of test results. Because these tests are conducted just prior to planting, their ability to forecast seedling growth on a reforestation site has a limited time frame. Consequently, a number of studies have reported various levels of success in forecasting growth on a reforestation site (Grossnickle et al. 1991a-c; Grossnickle and Major 1993 a,b; Major et al. 1993; Folk et al. 1993).


Inconsistencies in forecasting seedling growth in the field are due to several factors. First, errors in describing potential seedling performance can occur in a system that aggregates many plant physiological and morphological characteristics (e.g., cells, tissues, and organs) having different turnover times (Gardner et al. 1982). Seedlings have a dynamic pattern to their seasonal physiological response and morphological devel-



**Figure 1** - Possible testing procedures for determining seedling field performance potential in response to cold reforestation site environmental conditions.

opment (Fuchigami et al. 1982, Burr 1990, Ritchie and Tanaka 1990). Any testing procedure is just a "snapshot" of a single point in time along this seasonal pattern, making it difficult to accurately forecast all future seasonal patterns. Second, seedling field site performance may not always match stock quality test results because it is difficult to simulate all possible combinations of environmental stress—that is, duration, timing, intensity, frequency—that could occur under actual field site conditions. This makes it difficult to always define the proper level of environmental stress needed to obtain useful information on field performance potential that would forecast growth of seedlings on reforestation sites.

This does not mean that forecasting seedling field performance potential is not possible. One could come closer to defining a seedling's actual field response by using a greater number of material and performance attribute tests designed to give information on a seedling's overall response to potentially limiting site related environmental conditions. Also, information on typical seasonal trends of environmental conditions, for reforestation sites within defined ecosystems, could be used to develop test environments that provide a fair representation of what seedlings might



**Figure 2** - Possible testing procedures for determining seedling field performance potential in response to drought reforestation site environmental conditions.

be exposed to in the field. With this information, attributes such as those in table 1 could be selected to characterize a seedling's response to expected environmental conditions of a specific planting site.

In the following example, we describe how actual field response was forecasted by using a combination of material and performance attribute tests. Field performance potential was measured, under controlled laboratory conditions, on western hemlock (*Tsuga heterophylla* (Raf.) Sarg.) seedlings destined for late winter planting when exposure to low temperature conditions was probable (Grossnickle et al. 1991a). Western hemlock seedlings treated with short-day (compared to long-day) dormancy induction treatments had better field performance potential in the following tests (table 1, figure 1):

1. Water movement capability through the plant atmosphere continuum at 5 °C root temperature.
2. Net photosynthesis 14-day integral with root temperature at 5 °C.
3. Root growth capacity at a root temperature of 5 °C.

4. Frost hardiness of the whole shoot system to -18 /C.
5. Seedling water balance ratio.

One month after planting on a reforestation site, and after exposure to low temperatures and frosts in late winter and early spring, short-day treated seedlings had the least needle damage due to frosts and the greatest amount of new root growth (Grossnickle et al. 1991b). In addition, short-day treated seedlings had greater needle conductance and net photosynthesis after frost events during this late winter and early spring period (Grossnickle and Arnott 1992). In this example, material and performance attribute tests were selected in anticipation of low-temperature site conditions just after planting. This group of tests yielded a fairly accurate forecast of subsequent field performance.

Attributes defined in table 1 are not an all-inclusive list, but an example of parameters to consider for a comprehensive stock quality assessment program. Inclusion of alternative material or performance attribute tests in the master table is possible depending upon the user's needs and further development of testing procedures. A number of authors have identified additional seedling physiological and morphological characteristics that might be important for

inclusion in a master table (Timmis 1980, table 1; Burdett 1983, table 1; Puttonen 1989, table 2). Material and performance attribute tests need to be developed with these physiological and morphological characteristics in mind.

## Conclusions

Stock quality testing procedures that measure the functional integrity of seedlings at time of planting help to forecast initial survival capability. A number of

testing approaches are available that determine whether stock meets certain minimum physiological standards, thus allowing for removal of seedlings with poor field survival capability.

On the other hand, testing for field performance potential requires a separate approach to forecast seedlings physiological response and growth on a reforestation site. A combination of material and performance attributes tests, designed to provide information on a seedling's physiological response and growth to potentially limiting site-related environmental conditions, comes closer to defining actual field performance. Whether this approach is practical in all operational reforestation programs is questionable

The sophisticated equipment and technical expertise required to conduct field performance potential testing, as has been described, will limit its use. One can speculate that field performance potential testing could be beneficial to nursery personnel in developing new stock types or nursery cultural regimes. Regeneration silviculturists could use field performance potential testing when planting seedlings on field sites where survival and/or growth is known to be limited. Field performance potential testing has been used in our lab to test seedlings from a number of operational reforestation programs where field site conditions or stock type performance was considered limiting to reforestation success.

Stock quality testing using the above described approaches would provide a means for nursery personnel and regeneration silviculturists to better forecast initial field survival capability or field performance potential of seedlings. With this information, forest regeneration programs can work towards producing seedlings that meet the definition of stock quality—"fitness for purpose."

## Literature Cited

- Armson, K. A.; Sadreka, V. 1979. Forest tree nursery soil management and related practices. Toronto: Ontario Ministry of Natural Resources. 177 p.
- Binder, W D.; Scagel, R. K.; Krumlik, G. J. 1988. Root growth potential: Facts, myths, values? Gen. Tech. Rep. RM-167 Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station: 111-118.
- Brissette, J. C.; Chambers, J. L. 1992. Leaf water status and root system water flux in shortleaf pine (*Pinus echinata* Mill.) seedlings in relationship to new root growth after transplanting. *Tree Physiology* 11:289-303.
- Burdett, A. N. 1983. Quality control in the production of forest planting stock. *Forestry Chronicle* 59:132-138.
- Burdett, A. N. 1987. Understanding root growth capacity: Theoretical considerations in assessing planting stock quality by means of root growth tests. *Canadian Journal of Forest Research* 17:768-775.
- Burdett, A. N. 1990. Physiological processes in plantation establishment and the development of specifications for forest planting stock. *Canadian Journal of Forest Research* 20:415-427.
- Burr, K. E. 1990. The target seedling concept: Bud dormancy and cold-hardiness. In: Rose, R.; Campbell, S. J.; Landis, T D., eds. Target Seedling Symposium; Proceedings, Combined Meeting of the western Forest Nursery Associations. Gen. Tech. Rep. RM 200. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station: 79-90.
- Carlson, W C.; Miller, D. E. 1990. Target seedling root system size, hydraulic conductivity, and water use during seedling establishment. In: Rose, R.; Campbell, S. J.; Landis, T D., eds. Target Seedling Symposium: Proceedings, Combined Meeting of the Western Forest Nursery Associations. Gen. Tech. Rep. RM-200. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station: 53-66.

- Chavasse, C. G. R. 1980. Planting stock quality: A review of factors affecting performance. *New Zealand Journal of Forest Research* 25:144-171.
- Cleary, B. D.; Greaves, R. D.; Owston, P. W. 1978. Seedlings. In: Cleary, B. D.; Greaves, R. D.; Hermann, R. K., eds. *Regenerating Oregon forests*. Corvallis, OR: Oregon State University Extension Service: 63-98.
- Colombo, S. J. 1986. Second-year shoot development in black spruce *Picea mariana* (Mill.) B.S.P container seedlings. *Canadian Journal of Forest Research* 16:68-73.
- D'Aoust, A. D.; Delisle, C.; Girouard, R.; Gonzalez, A.; Bernier-Cardou, M. 1991. Characterization of containerized spruce seedlings prior to outplanting. In: Donnelly F. P.; Lussenburg, H. W., eds. *Proceedings, 11th Annual Conference of the Forest Nursery Association of British Columbia*. 59-63.
- DeLucia, E. H. 1987. The effect of freezing nights on photosynthesis, stomatal conductance, and internal CO<sub>2</sub> concentration in seedlings of Engelmann spruce (*Picea engelmannii* Parry). *Plant and Cell Environment* 10:333-338.
- Dixon, R. K.; Pallardy, S. G.; Garrett, H. E.; Cox, G. S. 1983. Comparative water relations of container-grown and bare-root ectomycorrhizal and nonmycorrhizal *Quercus velutina* seedlings. *Canadian Journal of Botany* 61:1559-1565.
- Duryea, M. L., ed. 1985a. Evaluating seedling quality: Principles, procedures, and predictive abilities of major tests. Corvallis, OR: Oregon State University, Forestry Research Laboratory. 143 p.
- Duryea, M. L. 1985b. Evaluating seedling quality: Importance to reforestation. In: Duryea, M. L., ed. *Evaluating seedling quality: Principles, procedures, and predictive abilities of major tests*. Corvallis, OR: Oregon State University, Forestry Research Laboratory: 1-6.
- Folk, R. S.; Grossnickle, S. C.; Major, J. E.; Arnott, J. T. 1993. Influence of nursery culture on western red cedar: 2. Freezing tolerance and morphological development of fall- and spring-planted seedlings. *New Forests* (in press).
- Fuchigami, L. H.; Weiser, C. J.; Kobayashi, K.; Timmis, R.; Gusta, L. V. 1982. A degree growth stage (°GS) model and cold acclimation in temperate woody plants. In: Li, P. H.; Sakai, A., eds. *Plant cold hardiness and freezing stress*. New York: Academic Press: 93-116.
- Gardner, R. H.; Cale, W. G.; O'Neill, R. V. 1982. Robust analysis of aggregation error. *Ecology* 63:1771-1779.
- Gerum, C. 1985. Frost hardiness of coniferous seedlings: Principles and applications. In: Duryea, M. L., ed. *Evaluating seedling quality: Principles, procedures, and predictive abilities of major tests*. Corvallis, OR: Oregon State University, Forestry Research Laboratory: 107-123.
- Gerum, C. 1988. Evaluation of planting stock quality. In: Smith, C. R.; Reffle, R. J., eds. *Taking stock: The role of nursery practice in forest renewal*. OFRC Symposium Proc. O-P-16. Sault Ste. Marie, ON: Canadian Forestry Service, Great Lakes Forestry Centre: 44-49.
- Grossnickle, S. C. 1988. Planting stress in newly planted jack pine and white spruce. 1. Factors influencing water uptake. *Tree Physiology* 4:71-83.
- Grossnickle, S. C.; Arnott, J. T. 1992. Gas exchange response of western hemlock seedlings from various dormancy induction treatments to reforestation site environmental conditions. *Forest Ecology and Management* 49:177-193.
- Grossnickle, S. C.; Blake, T. J. 1987. Comparison of water relation patterns for newly planted bare-root and container jack pine and black spruce seedlings on boreal cut-over sites. *New Forests* 1:101-116.
- Grossnickle, S. C.; Heikurinen, J. 1989. Site preparation: Water relations and growth of outplanted jack pine and white spruce. *New Forests* 3:99-123.
- Grossnickle, S. C.; Major, J. E. 1993a. Interior spruce seedlings compared to emblings produced from somatic embryogenesis: 2. Stock quality assessment prior to field planting. (unpublished article)
- Grossnickle, S. C.; Major, J. E. 1993b. Interior spruce seedlings compared to emblings produced from somatic embryogenesis: 3. Physiological response and morphological development on a reforestation site. (unpublished article)
- Grossnickle, S. C.; Reid, C. P. P. 1984. Water relations of Engelmann spruce seedlings on a high-elevation mine site: An example of how reclamation techniques can alter microclimate and edaphic conditions. *Reclamation and Revegetation Research* 3:199-221.
- Grossnickle, S. C.; Arnott, J. T.; Major, J. E. 1988. A stock quality assessment procedure for characterizing nursery-grown seedlings. In: *Proceedings, Combined Nursery Meetings of the Western Forest Nursery Associations*. Gen. Tech. Rep. RM-167 Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station: 77-88.
- Grossnickle, S. C.; Arnott, J. T.; Major, J. E.; Tschaplinski, T. J. 1991a. Influence of dormancy induction treatments on western hemlock seedlings. 1. Seedling development and stock quality assessment. *Canadian Journal of Forest Research* 21:164-174.
- Grossnickle, S. C.; Arnott, J. T.; Major, J. E.; LeMay, V. M. 1991c. Stock quality assessment through an integrated approach. *New Forests* 5:77-91.
- Hawkins, C. B. D.; Binder, W. D. 1990. State of the art stock quality tests based on seedling physiology. In: Rose, R.; Campbell, S. J.; Landis, T. D. *Target Seedling Symposium: Proceedings, Combined Meeting of the Western Forest Nursery Associations*. Gen. Tech. Rep. RM-200. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station: 91-122.
- Hawkins, C. B. D.; Deyoe, D. R. 1992. SIVE, a new stock quality test: The first approximation. *FRDA Rep. 175*. Victoria, BC: British Columbia Ministry of Forestry and Forestry Canada. 24 p.
- Hinckley, T. M.; Lassoie, J. P.; Running, S. W. 1978. Temporal and spatial variations in the water status of forest trees. *Forestry Sciences Monograph* 20. Washington, DC: Society of American Foresters. 72 p.
- Jaramillo, A. 1981. Review of techniques used to evaluate seedling quality. Gen. Tech. Rep. INT-109. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experiment Station: 84-95.
- Johnson, J. D.; Cline, M. L. 1991. Seedling quality of southern pines. In: Duryea, M. L.; Dougherty P. M. *Forest regeneration manual*. Dordrecht, The Netherlands: Mower Academic Publishers: 143-162.
- Kaufmann, M. R. 1977. Soil temperature and drying cycle effects on water relations of *Pinus radiata*. *Canadian Journal of Botany* 55:2413-2418.
- Kaufmann, M. R. 1979. Stomatal control and the development of water deficit in Engelmann spruce seedlings during drought. *Canadian Journal of Forest Research* 9:297-304.
- Kozlowski, T. T.; Davies, W. J. 1975. Control of water balance in transplanted trees. *Arboretum* 1:1-10.
- Landseberg, J. J. 1986. *Physiological ecology of forest production*. New York: Academic Press.
- Langerud, B. R. 1991. "Planting stock quality": A proposal for better terminology. *Scandinavian Journal of Forest Research* 6:49-51.
- Langerud, B. R.; Puttonen, P.; Troeng, E. 1991. Viability of *Picea abies* seedlings with damaged roots and shoots. *Scandinavian Journal of Forest Research* 6:59-72.

- Lavender, D. P. 1988. Characterization and manipulation of the physiological quality of planting stock. In: Worrall, J.; Loo-Dinkins, J.; Lester, D., eds. Proceedings, 10th North American Forest Biology Workshop, Physiology and Genetics of Reforestation. Vancouver, BC: University of British Columbia: 32-57.
- Lavender, D. P. 1991. Measuring phenology and dormancy. In: Lassoie, J. P.; Hinckley T M., eds. Techniques and approaches in forest tree ecophysiology. Boca Raton, FL: CRC Press, 403-422.
- Levitt, J. 1980. Responses of plants to environmental stresses: vol. 1, Chilling, freezing, and high temperature stresses, 2nd ed. New York: Academic Press.
- Livingston, N. J.; Black, T. A. 1987a. Water stress and survival of three species of conifer seedlings planted on a high elevation south-facing clear-cut. Canadian Journal of Forest Research 17:1115-1123.
- Livingston, N. J.; Black, T. A. 1987b. Stomatal characteristics and transpiration of three species of conifer seedlings planted on a high elevation south-facing clear-cut. Canadian Journal of Forest Research 17:1273-1282.
- Livingston, N. J.; Black, T. A. 1988. The growth and water use of three species of conifer seedlings planted on a high elevation south-facing clear-cut. Canadian Journal of Forest Research 18:1234-1242.
- Lopushinsky, W.; Kaufmann, M. R. 1984. Effects of cold soil on water relations and spring growth of douglas-fir seedlings. Forest Science 30:628-634.
- Major, J. E.; Grossnickle, S. C.; Folk, R. S.; Arnott, J. T. 1993. Influence of nursery culture on western red cedar: 1. Measurement of seedling attributes before fall and spring planting. New Forests (in press).
- Margolis, H. A.; Waring, W. H. 1986. Carbon and nitrogen allocation patterns of Douglas-fir seedlings fertilized with nitrogen in the autumn. 2. Field performance. Canadian Journal of Forest Research 16:903-909.
- McCreary, D. D.; Duryea, M. L. 1985. OSU vigor test: Principles, procedures, and predictive ability. In: Duryea, M. L., ed. Evaluating seedling quality: Principles, procedures, and predictive ability of major tests. Corvallis, OR: Oregon State University Forestry Research Laboratory: 85-92.
- McCreary, D. D.; Duryea, M. L. 1987. Predicting field performance of Douglas-fir seedlings: Comparison of root growth potential, vigor, and plant moisture stress. New Forests 2:153-169.
- McKay, H. M. 1992. Electrolyte leakage from fine roots of conifer seedlings: A rapid index for plant vitality following cold storage. Canadian Journal of Forestry Research 22:1371-1377.
- McKay H. M.; Mason, W. L. 1991. Physiological indicators of tolerance to cold storage in Sitka spruce and Douglas-fir seedlings. Canadian Journal of Forestry Research 21:890-910.
- Miller, P. C. 1983. Comparison of water balance characteristics of plant species in "natural" versus modified ecosystems. In: Mooney, H. A.; Gordon, M. Disturbance and ecosystems: Components of response. New York: Springer-Verlag: 188-212.
- Nambiar, E. K. S.; Zed, P. G. 1980. Influence of weeds on the water potential, nutrient content and growth of young radiata pine. Australian Forestry Research 10:279-288.
- Nambiar, E. K. S.; Bowen, G. D.; Sands, R. 1979. Root regeneration and plant water status of *Pinus radiata* D. Don seedlings transplanted to different soil temperatures. Journal of Experimental Botany 30:1119-1131.
- Neilson, R. E.; Jarvis, P. G. 1976. Photosynthesis in Sitka spruce (*Picea sitchensis* (Bong.) Carr.): 6. Response of stomata to temperature. Journal of Applied Ecology 12:879-891.
- Nilsson, J.; Eriksson, G. 1986. Freeze testing and field mortality of *Pinus sylvestris* (L.) in northern Sweden. Scandinavian Journal of Forest Research 1:205-218.
- Omi, S. K. 1991. The target seedling and how to produce it. In: van Buijtenen, J. P.; Simms, T., eds. Nursery management workshop. Pub. 148. College Station, TX; Texas Forest Service: 88-118.
- Orlander, G.; Rosvall-Ahnebrink, G. 1987. Evaluating seedling quality by determining their water status. Scandinavian Journal of Forest Research 2:167-177.
- Orlander, G.; Egnell, G.; Forsen, S. 1989. Infrared thermography as a means of assessing seedling quality. Scandinavian Journal of Forest Research 4:215-222.
- Puttonen, P. 1989. Criteria for using seedling performance potential tests. New Forests 3:67-87.
- Ritchie, G. A. 1984. Assessing seedling quality. In: Duryea, M. L.; Landis, T. D., eds. Forest nursery manual: Production of bareroot seedlings. The Hague: Martinus Nijhoff/Dr. W Junk Publishers: 243-259.
- Ritchie, G. A. 1985. Root growth potential: Principles, procedures, and predictive ability. In: Duryea, M. L., ed. Evaluating seedling quality: Principles, procedures, and predictive ability of major tests. Corvallis, OR: Oregon State University, Forestry Research Laboratory: 93-106.
- Ritchie, G. A. 1990. A rapid method for detecting cold injury in conifer seedling root systems. Canadian Journal of Forest Research 20:26-30.
- Ritchie, G. A.; Dunlap, J. R. 1980. Root growth potential: Its development and expression in forest tree seedlings. New Zealand journal of Forest Science 10:218-248.
- Ritchie, G. A.; Tanaka, Y. 1990. Root growth potential and the target seedling. In: Rose, R.; Campbell, S. J.; Landis, T. J., eds. Target Seedling Symposium, Proceedings, Combined Meeting of the Western Forest Nursery Associations. Gen. Tech. Rep. RM-200. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station: 37-51.
- Sands, R. 1984. Transplanting stress in radiata pine. Australian Forest Research 14:67-72.
- Schmidt-Vogt, H. 1981. Morphological and physiological characteristics of planting stock: Present state of research and research tasks for the future. In: Proceedings, IUFRO 17th World Congress; Tokyo: 433-446.
- Stone, E. C. 1955. Poor survival and the physiological condition of planting stock. Forest Science 1:89-94.
- Sutton, R. E. 1979. Planting quality and grading. Forest Ecology and Management 2:123-132.
- Sutton, R. P. 1982. Plantation establishment with bareroot stock: Some critical factors. In: Mroz, G. D.; Berner, J. E., eds. Artificial regeneration of conifers in the upper Great Lakes region. Houghton, MI: Michigan Technological University: 304-331.
- Sutton, R. F. 1988. Planting stock quality is fitness for purpose. In: Smith, C. R.; Reffle, R. J., eds. Taking stock: The role of nursery practice in forest renewal. Proceedings, OFRC symposium. O-P 15. Sault Ste. Marie, ON: Canadian Forestry Service, Great Lakes Forestry Centre: 39-43.
- Thompson, B. E. 1985. Seedling morphological evaluation: What you can tell by looking. In: Duryea, M. L., ed. Evaluating seedling quality: Principles, procedures, and predictive ability of major tests. Corvallis, OR: Oregon State University, Forestry Research Laboratory: 59-72.
- Timmis, R. 1980. Stress resistance and quality criteria for tree seedlings: Analysis, measurement, and use. New Zealand journal of Forest Science 10:21-53.
- Tyree, M. T.; Jarvis, P. G. 1982. Water in tissues and cells. In: Lange, O. L.; Nobel, P. S.; Osmond, C. B.; Ziegler, H. Physiological plant ecology. Vol. 2. Water relations and carbon assimilation. New York: Springer-Verlag: 36-77.

- Vanhinsberg, N. B.; Colombo, S. J. 1990. Effect of temperature on needle anatomy and transportation of *Picea mariana* after bud initiation. Canadian Journal of Forest Research 20:598-601.
- Vidaver, N.; Binder, W; Brooke, R. C.; Lister, G. R.; Toivonen, P M. A. 1989. Assessment of photosynthetic activity of nursery-grown *Picea glauca* seedlings using an integrating fluorometer to monitor variable chlorophyll fluorescence. Canadian Journal of Forest Research 19:1478-1482.
- Vidaver, N.; Lister, G. R.; Brooke, R. C.; Binder, W 1991. A manual for the use of variable chlorophyll fluorescence in the assessment of the ecophysiology of conifer seedlings. FRDA Rep. 163. Victoria, BC: British Columbia Ministry of Forests and Forestry Canada. 166 p.
- Weatherspoon, P; Laacke, J. 1985. Infrared thermography for assessing seedling condition: Rational and preliminary observations. In: Duryea, M. L., ed. Evaluating seedling quality: Principles, procedures, and predictive ability of major tests. Corvallis, OR: Oregon State University, Forestry Research Laboratory: 127-136.