

## Materials and Methods

Italian alder and Douglas-fir seeds were collected, respectively, in 1991 in Tivoli, Rome (41,58 N; 12,48 E; 50 m asl) and in 1980 in Albany, Oregon (44,38 N; 123,06 W; 700 m asl). They were stored at -3 /C (27 /F).

For Italian alder, germination percentage was 48%; purity, 89%; moisture content, 7%; and the number of cleaned seeds per kilogram, 540,000. For Douglas-fir, germination percentage was 84.5%; moisture content, 8.5%; and the number of cleaned seeds per kilogram, 88,500. Seeds for trials were obtained from four 50-g (1.7-oz) samples withdrawn at random from both seedlots.



The DWD composition was sucrose, 50%; betaine, 20%; valine, 10%; isoleucine, 5%; leucine, 3%; tyrosine, 10%; glycine, glutamic acid, and serine, trace; other components, 2%. For each species, a 2-factor split-plot (Gomez and Gomez 1984) design with 4 replications was used in the experiment. There were 2 pregermination treatments; (A1) prechilling at 4 /C (39 /F) for 21 days and (A2) no prechilling. Three concentrations of DWD were tested: (B1) Control (0 ppm); (B2) 10 ppm; (B3) 30 ppm. The 6 treatments resulting from treatment combinations are shown in table 1.

During the chilling period, seeds were placed in germination boxes with water. They were then rinsed in water and put in new boxes containing DWD solutions at different concentration (0, 10, or 30 ppm) for the germination test. Unchilled seeds were soaked in water for 24 hours at room temperature and then placed in germination boxes containing DWD solutions (0, 10 or 30 ppm). Prechilling was performed so that all seeds could begin germination tests at the same time in a cabinet germinator. In both species, four 50-seed replicates were prepared for each of the 6 experimental treatments (table 1). Seeds were placed on top of filter paper in covered germination boxes (diameter = 16 cm) in which water or solutions uptake was easily allowed through strips of filter paper from a reservoir at the bottom (figure 1A&B). Germination boxes contained 250 ml of water or DWD solutions, depending on experimental treatment.

Germination conditions were those prescribed by the ISTA (1993) rules: 8 hours at 30 /C (86 /F) in light plus 16 hours at 20 /C (68 /F) in dark. Experimental treatments A2B1 for Italian alder and A1B1 and A2B1 for Douglasfir are to be considered under the standard conditions as they followed ISTA (1993) recommended pregermination treatments: no prechilling for Italian alder, double test (prechilling and no prechilling) for these 2 species as already stated. Germination was recorded when the radicle length exceeded 2 mm (Danielson and Tanaka

**Table 1**—Prechilling (factor A) and distillary wash derivative (DWD) (factor B) treatments applied to Italian alder and Douglas-fir seeds

Pregermination treatment	DWD conc. (ppm)
Prechilling (A1)	0—control (B1)
	10 (B2)
	30 (B3)
No prechilling (A2)	0—control (B1)
	10 (B2)
	30 (B3)



**Figure 1**—Germination box; seen from the top (A); detail of the box showing uptake water or solutions system (B).



1978; De Matos Malavasi and others 1985). The germinated seeds were counted every 2 or 3 days until the 28th day to calculate the germination percentage (G%) on the 3rd, 7th, and 28th day (final germination) and the mean time to complete germination (MTG) (Bewley and Black 1986). MTG is a measure of the speed of germination:

$$3(t \times n)/3n$$

where:  $t$  = the time in days, starting from the day of the beginning of the test  
 $n$  = the number of seeds completing germination on day  $t$

Quicker germination is associated with lower MTG values (Bewley and Black 1986).

Statistical differences were estimated on the basis of the analysis of variance (Gomez and Gomez 1984) of MTG's and G% at arcsin % (tabulated values are the untransformed data). Comparison of means was performed following the Duncan's multiple range test (Harter 1960) at  $P < 0.01$ .

## Results and Discussion

Data presented in tables 2 and 3 show that statistically significant differences were only attributable to main factors. In Italian alder, MTG was more influenced than percentage germination, with this effect being more noticeable at the beginning than at the end of the germination process. Rapid germination processes are often associated with remarkable early G%'s. In Douglas-fir, both cold treatment and DWD treatment affected speed and total germination.

**Pregermination treatments (factor A).** In both species, seeds that were not subjected to prechilling (A2; showed a low G% at the beginning of the trial (3rd day in Italian alder and 7th day in Douglas-fir). Final germination (28th day) of pretreated Italian alder seeds did not significantly differ from those that were stratified (A1). In Douglas-fir, cold-moist pretreatment resulted in higher early and final germination of prechilled seeds in comparison to untreated seeds (table 2).

Germination speed, expressed as MTG, was always improved by stratification (table 2). On the 3rd day, G%, was equal to almost 70% of final germination percentage in stratified Italian alder seeds (A1), whereas no germination could be observed in untreated seeds. Such an effect was expected in both species because it shows consistency with widespread nursery practice in which both Italian alder or Douglas-fir seeds are subject to stratification before spring sowing (Gordon and Rowe

1982). Although this trial was conducted with only 1 provenance, the results suggest a probable need of modification of ISTA rules, that prescribe prechilling before the germination test for Italian alder.

### DWD concentration during germination test (factor B).

The use of DWD solutions during germination did not affect the rate or the completeness of germination in Douglas-fir but it did accelerate the germination process in Italian alder (table 3). Furthermore, DWD solutions increased early G% (3rd day), and thus germination speed, in the latter species. There was no difference between 10 and 30 ppm (table 3). It was hypothesized that the combination of sucrose and amino acids present in DWD might have resulted in rapid embryo development in Italian alder. Douglas-fir seeds were not affected by the DWD concentrations tested, probably due to their thicker seedcoat.

## Conclusions

Results of the study have indicated that:

- The use of DWD solutions (10 and 30 ppm) during germination tests accelerated the germination process in Italian alder but did not affect the rate or the completeness of germination in Douglas-fir. We hypothesize that differences in seed coat structure could explain different behaviours and responses.
- Stratification strongly influenced both early G% and MTG in Italian alder and early, final G% and MTG in Douglas-fir. Although prechilling is prescribed by ISTA (1993) rules for germination tests in the former species it is not indicated for Italian alder. These results could suggest a probable need of modification of ISTA rules.

Although preliminary in nature, the results show that DWD improved tree seed germination in 1 species. Further research is required on a wider range of species, as DWD is cheap, safe, and easy to use.

**Address correspondence to:** Dr. Elvira Rea, Istituto, Sperimentale per la Nutrizione delle Plante, Via delta Navicella 2-4, 00184, Roma, ITALY; **e-mail:** » [fisveg@www.inea.it](mailto:fisveg@www.inea.it) « Dr. Piotta is currently affiliated with the Ministero dell'Ambiente SIAR, Via della Ferratella in

**Table 2**—Germination percentages on the 3rd, 7th, and 28th days and mean time to complete germination (MTG) of seeds of Italian alder and Douglas-fir subjected to different pregermination treatments

	Italian alder		Douglas-fir	
	Prechilling	No prechilling	Prechilling	No prechilling
Germination (%)				
3rd day	35.0 B b	0.01 A a	ng	ng
7th day	46.9 A a	51.9 A a	35.9 B b	4.2 A a
28th day	48.2 A a	57.0 A b	73.5 B b	67.5 A a
MTG (days)	3.9 A	6.3 B	9.2 A a	14.8 B b

**Note:** ng = no germination observed. Values are averages for 3 distillery wash derivative (DWD) solutions. For each species, means within a row followed by different capital letters are significantly different at  $P < 0.01$  according to the Duncan's multiple range test (Harter 1960); means within a row followed by different lowercase letters are significantly different at  $P < 0.05$ .

**Table 3**—Germination percentages on the 3rd, 7th, and 28th day and mean time to complete germination (MTG) of seeds of Italian alder and Douglas-fir imbibed with distillery wash derivative (DWD) solutions of 0, 10, & 30 ppm during the germination test

	Italian alder			Douglas-fir		
	0 ppm	10 ppm	30 ppm	0 ppm	10 ppm	30 ppm
Germination (%)						
3rd day	6.9 A a	12.8 B b	10.7 AB b	ng	ng	ng
7th day	52.6 A a	49.3 A a	47.8 A a	16.7 A a	14.6 A a	19.7 A a
28th day	56.3 A a	52.4 A a	49.3 A a	66.6 A a	70.7 A a	75.0 A a
MTG (days)	5.6 b B b	4.9 AB a	4.8 A a	12.4 A b	12.0 A ab	11.5 A a

**Note:** ng = no germination observed. Values are averages for 2 pregermination treatments. For each species, means within a row followed by different capital letters are significantly different at  $P < 0.01$  according to the Duncan's multiple range test (Harter 1960); means within a row followed by different small letters are significantly different at  $P < 0.05$ .

## References

- Bewley JD, Black M. 1986. Seeds, physiology of development and germination. New York: Plenum Press. 4 p.
- Bonner FT, McLemore BF, Barnett JP 1974. Presowing treatment of seed to speed germination. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk 450. Washington, DC: USDA Forest Service: 126-135.
- Cullum FJ, Cordon AG. 1993. Dormancy release of tree and shrub seeds using a compost activator pretreatment. Proceedings of the International Plant Propagators' Society 43:125-130.
- Danielson HR, Tanaka Y. 1978. Drying and storing stratified ponderosa pine and Douglas fir seeds. Forest Science 24(1): 11-16.
- De Matos Malavasi M, Stafford SG, Lavender DP 1985. Stratifying, partially redrying and storing Douglas-fir seeds: effects on growth and physiology during germination. Annales des Sciences Forestieres 42(4): 371-384.
- Di Monte G, Bragaloni M, Pierandrei F, Rea E. 1995. Effect of distillery washes on seed germination. Giornale Botanico Italiano 129(4): 999-1000.
- Gomez KA, Gomez AA. 1984. Statistical procedures for agricultural research. New York: John Wiley & Sons: 97-107.
- Cordon AE. 1979. The effect of seed maturity on the dormancy and the viability of British produced Douglas-fir and Sitka spruce seeds. In: Bonner F, ed. Proceedings, Symposium on Flowering and Seed Development in Trees; 1979. Starkville, MS: USDA Forest Service, Southern Forest Experiment Station: 362.
- Cordon AE, Rowe DC. 1982. Seed manual for ornamental trees and shrubs. Forestry Commission Bulletin 59: 120-128.
- Halmer P. 1989. The use of plant growth regulators in the production and treatment of seeds: a commercial prospective. In: Pinfield NJ, Back M, eds. British Plant Growth Regulator Group Monograph 15: 77-92.
- Harter HL. 1960. Critical values for Duncan's new multiple range test. Biometrics 16: 675-685.
- Heydecker W, Coolbear P 1977. Seed treatments for improved performance: survey and attempted prognosis. Seed Science and Technology 5: 353-425.
- ISTA [International Seed Testing Association]. 1993. International rules for seed testing 1993. Seed Science and Technology 21 (suppl.): 160-186.
- Oplinger ES, Anderson IR, Johnson RR. 1978. Effect of seed and foliar applications of Ergostim on soybeans and corn. In: Abdel Rahman M, ed. Proceedings, 5th Annual Meeting of the Plant Growth Regulator Working Group (PGRWG); Blacksburg, VA; 1978 June 25-29: 124-128.
- Orsi S, Tallarico R. 1983. Fitoregolatori a biostimolanti su grano ed orzo. L'Informatore Agrario 38: 539-541.

- Owston PW, Stein WI. 1974. *Pseudotsuga* Carr., Douglas-fir. In: Schopmeyer CS, tech. word. Seeds of woody plants in the United States. Agric. Handbk. 450: 674-680.
- Piotto B. 1992. Semi di alberi a arbusti coltivati in Italia, come e quando seminarli. Rome: SAF [Centro di Sperimentazione Agricola a Forestale]: 78.
- Rinallo C. 1979. Alcuni aspetti della biologia dello sviluppo in ontano napoletano (*Alnus cordata* Loisel). Annali dell'Accademia Italiana di Scienze Forestali 28: 207-226.
- Siviero P 1993. Effetti di applicazioni di biostimolanti su pomodoro a melone. L'Informatore Agrario 45: 55-60.
- Vanesse R. 1974. Evolution du degre de dormance des graines de sapin de Douglas (*Pseudotsuga menziesii* (Mirb.) Franco). Bulletin de la Societe Royale Forestiere de Belgique 81(3): 149-158.